

What is Malnutrition?

Malnutrition is the condition that develops when the body does not get the right amount of vitamins, minerals, and other nutrients (e.g. energy, protein) it needs to maintain health, promote cell and tissue growth and normal organ function. Malnutrition may result from consuming too little food, a shortage of key nutrients, or altered absorption or metabolism. Older adults are at particular risk of malnutrition.



The financial costs associated with malnutrition are huge. It is estimated that the cost of malnutrition to the EU alone is a staggering €170 billion.¹

Malnutrition: A Rising Issue


The World Health Organisation (WHO) estimates that by 2015, malnutrition will affect 1 in 6 of the global population.² In Europe alone the issue of malnutrition impacts more than 30 million citizens.

- Malnutrition can be found in all healthcare settings. Hospitalised patients are at particular risk as 86% of them have been identified as malnourished or at risk of malnutrition³
- Malnutrition is expected to become an even greater problem as a result of an ageing population and an increase in chronic diseases that are often associated with malnutrition (e.g. dementia)⁴
- Despite its high prevalence, physician awareness of the important role nutrition plays in general well being and disease treatment is quite low. This results in a delay or omittance of appropriate nutrition intervention and leaves many people suffering the consequences of malnutrition⁵

In the older adult population:

- 50% eat less than the RDA for protein⁶
- 90% are Vitamin D deficient⁷
- 30% are Vitamin B₁₂ deficient⁸
- 30% have inadequate Zinc and Selenium intake^{9,10}

Prevalence of Malnutrition by Healthcare Setting³



Causes of Malnutrition

Older persons are particularly vulnerable to malnutrition. The process of aging affects nutrient needs – some nutrient requirements increase while others decrease. This often translates to the need for more nutrient dense food sources – allowing one to receive the needed nutrients in less food volume. Decrease of appetite, dental problems, psychosocial issues, illness and chronic disease often result in lower energy intake and lower intakes of essential nutrients. Recent data from the European Nutrition Day study showed that less than 40% of patients eat all the food they are served in the hospital.¹¹

Malnutrition Impairs Outcome

Malnutrition has been shown to correlate with higher rates of mortality, longer length and increased cost of hospital stay.¹²⁻¹⁵

The presence of malnutrition puts individuals at risk of developing problems such as an increased risk of infection, delayed wound healing, impaired respiratory function, muscle weakness, falls, fractures and delayed recovery.


Malnourished patients have a:

- 2 fold increased risk of long-term mortality¹⁵
- 3 times longer length of hospitalisation^{16,17}
- 3 times higher risk of infection¹³
- Higher costs of hospital care¹⁸
- Greater likelihood of hospital readmission after discharge¹⁹
- Greater dependence in activities of daily living (ADLs)


Low levels of vitamin E, B₁₂ and D have been associated with a decline in functional mobility

Malnourished patients have:


Up to 3 times higher risk of infection²⁰



A longer length of hospital stay¹⁶



Combined with disease, malnutrition puts patients at risk of entering a life-threatening, accelerated, downward spiral, potentially leading to dependence and institutionalisation.



Identifying Malnutrition

Malnutrition is an under-recognised and under-treated condition. This under-diagnosis may be attributed to inadequate education and training of healthcare professionals on the important role of nutrition in health and disease, and the inadequacy of nutrition screening programs. Screening and appropriate, timely nutritional intervention will help to reverse the negative consequences of malnutrition. Adequate nutritional support has been shown to shorten the length of hospital stay and improve quality of life both of which contribute to a reduction in the overall cost of care.^{21,22}

References:

1. Ljungqvist O. Presentation: The Cost of Malnutrition. June 11, 2009, Prague, CZ.
2. de Onis, M *et al.* (2004) Estimates of Global Prevalence of Childhood Underweight in 1990 and 2015. *The Journal of the American Medical Association*; **291**: 2600-2606.
3. Kaiser MJ *et al.* (2009) World-Wide Data on Malnutrition in the Elderly According to the Mini Nutritional Assessment® (MNA) – Insights from an International Pooled Database. *Clinical Nutrition*; **4** (S2): 113.
4. Elia, M *et al.* (2008) Combating Malnutrition: Recommendations for Action, Redditch, UK, British Association for Parenteral and Enteral Nutrition.
5. Norman, K *et al.* (2008) Prognostic impact of disease-related malnutrition. *Clinical Nutrition*; **27**: 5-15.
6. Kant AK *et al.* (1999) Relation of Age and Self-reported Chronic Medical Condition Status with Dietary Nutrient Intake in the US Population. *J of Amer Coll Nutr*; **18**:69-76.
7. Cherniack EP *et al.* (2008) Hypovitaminosis D in the Elderly: From Bone to Brain. *J of Nutr Health and Aging*; **12**:366-373.
8. Bates CJ *et al.* (2002) Nutrition and aging: A consensus statement. *Jour of Nutr Health and Aging*; **6**:103-116.
9. Abellan van Kan G *et al.* (2008) Nutrition and Aging: The Carla Workshop. *Jour of Nutr Health and Aging*; **12**: 355-364.
10. Lauretani F *et al.* (2007) Association of low plasma selenium concentrations with poor muscle strength in older community-dwelling adults: the InCHIANTI Study. *Am J Clin Nutr*; **86**:347-352.
11. Hiesmayr M *et al.* (2009) Decreased food intake is a risk factor for mortality in hospitalised patients: NutriitionDay survey 2006. *Clinical Nutrition*; **28**:484 – 491.
12. Correia, ITD *et al.* (2003) Prevalence of Hospital Malnutrition in Latin America: *The Multicenter ELAN Study*. *Nutrition*; **19**: 823-825
13. Pirlich, M *et al.* (2006) The German hospital malnutrition study. *Clinical Nutrition*; **25**: 563-572.
14. Ockenga, J *et al.* (2005) Nutritional assessment and management in hospitalised patients: Implications for DG-based reimbursement and health care quality. *Clinical Nutrition*; **24**: 913-919
15. Sullivan DH *et al.* (2002) The GAIN (Geriatric Anorexia Nutrition) registry: the impact of appetite and weight on mortality in a long-term care population. *Jour of Nutr Health and Aging*; **6**: 275-281.
16. Pichard C *et al.* (2004) Nutritional Assessment : Lean body mass depletion at hospital admission is associated with an increased length of stay. *Am J Clin Nutr*; **79**:613-618.
17. Smith PE, Smith AE. (1997) High-quality nutritional interventions reduce costs. *Healthcare Finance Management*; **51**:66-69.
18. Chima CS *et al.* (1997) Relationship of nutritional status to length of stay, hospital costs, and discharge status of patients hospitalized in the medicine service. *J Amer Diet Assoc*; **97**: 979-80.
19. Thomas DL *et al.* (2002) Malnutrition in subacute care. *Am J Clin Nutr*; **75**:308-13.
20. Schneider SM *et al.* (2004) Malnutrition is an independent factor associated with nosocomial infections. *Br J Nutr*; **92**:105-11.
21. Lennard-Jones, J. (Chair) (1992) A Positive Approach to Nutrition as Treatment. London: Kings Fund Centre.
22. Green, C.J. (1999) Existence, causes and consequences of disease-related malnutrition in the hospital and the community, and clinical and financial benefits of nutritional intervention. *Clinical Nutrition*; **18**: Supp 2: 3-28.

®Société des Produits Nestlé S.A., Vevey, Switzerland, Trademark Owners.